Mean transverse mass of hadrons in proton-proton reactions

V. Yu. Vovchenko,[1,2,3] D. V. Anchishkin,[4,1] and M. I. Gorenstein[4,2]

[1] *Taras Shevchenko National University of Kiev, Kiev 03022, Ukraine*

[2] *Frankfurt Institute for Advanced Studies, Frankfurt 60438, Germany*

[3] *Goethe University, Frankfurt 60325, Germany*

[4] *Bogolyubov Institute for Theoretical Physics, Kiev 03680, Ukraine*

(Dated: January 26, 2015)

arXiv:1408.5493v2 [nucl-th] 23 Jan 2015

Abstract

An energy dependence of the mean transverse mass $\langle m_T \rangle$ at mid-rapidity in proton-proton ($p+p$) reactions is studied within the ultra-relativistic quantum molecular dynamics (UrQMD). The UrQMD model predicts a nonmonotonous dependence of $\langle m_T \rangle$ on collision energy for several hadron species: for π^+, p, K^+, and Λ the mean transverse mass has a maximum at the center of mass energy region $5 \leq \sqrt{s} \leq 8$ GeV. These results are a consequence of an interplay of two contributions: 1) excitations and decays of the baryonic resonances N^* and Δ; 2) excitations and decays of the baryonic strings. The UrQMD results do not show any nonmonotonous dependence of $\langle m_T \rangle$ on $\sqrt{s}$ for π^-, K^-, and antiprotons. Whether a nonmonotonous dependence of $\langle m_T \rangle$ at mid-rapidity on the collision energy for π^+, p, K^+, and Λ is relevant for real $p+p$ interactions will be soon checked experimentally by the NA61/SHINE Collaboration.

PACS numbers: 25.75.Gz, 25.75.Ag

Keywords: Proton-proton interactions; Mean transverse mass

Experimental data on hadron production in central Pb+Pb collisions obtained by the NA49 Collaboration [1–3] at the Super Proton Synchrotron (SPS) of the European Organization for Nuclear Research (CERN) are consistent with the onset of deconfinement in central nucleus-nucleus collisions at about $\sqrt{s_{NN}} = (7-8)$ GeV [4–6], where $\sqrt{s_{NN}}$ is the center of mass energy of two nucleons from colliding nuclei. An important experimental result in this context is an observation of a *step*-like behavior of a mean value of the transverse mass $m_T = \sqrt{m^2 + p_T^2}$, where m is a particle mass and p_T its transverse momentum. The mean transverse mass $\langle m_T \rangle$ for different hadron species (π^-, $K^\pm$, p, and $\overline{p}$) measured at mid-rapidity in central Pb+Pb collisions demonstrates an increase at small and large collision energies, but it remains approximately constant in the SPS energy range $\sqrt{s_{NN}} = 7.6-17.3$ GeV (see Fig. 8 in Ref. [3]). This is the energy region, where one expects the transition between confined and deconfined matter with the creation of mixed phase. Such an energy dependence of $\langle m_T \rangle$ is indeed typical for a 1^{st} order phase transition in the mixed-phase region [7–9]. In existing up to now data, there were no indications on such type of $\langle m_T \rangle$ dependence on the center of mass energy $\sqrt{s}$ in proton-proton ($p+p$) reactions.

A further progress in understanding the effects related to the onset of deconfinement can be achieved by a new comprehensive study of hadron production in $p+p$, proton-nucleus, and nucleus-nucleus collisions. This motivated the present NA61/SHINE ion programme at the SPS CERN devoted to the system size and energy scan [10, 11] and the Beam Energy Scan (BES) programme at the Relativistic Heavy Ion Collider (RHIC) of the Brookhaven National Laboratory (BNL) [12]. These efforts will be extended by the future Compressed Baryonic Matter (CBM) experiment at the Facility for Antiproton and Ion Research (FAIR) [13, 14], which will employ high-luminosity beams and large-acceptance detectors to study system-size and energy dependence of hadron production in proton-nucleus and nucleus-nucleus collisions.

In Ref. [15], we analyzed recent NA61/SHINE data [16] on π^- spectra in $p+p$ reactions within the ultra-relativistic quantum molecular dynamics (UrQMD) model [17, 18]. The version UrQMD-3.3p2 [19] was used in Ref. [15]. Recently a new version, UrQMD-3.4 [20], was released. It will be used in the present study. Both the data and the UrQMD simulations demonstrate a monotonous increase of $\langle m_T \rangle$ with collision energy for π^- at mid-rapidity. This is shown in Fig. 1 *left*. Unexpectedly, the UrQMD has predicted a nonmonotonous dependence of $\langle m_T \rangle$ on

collision energy for π^+ in $p+p$ reactions: the mean transverse mass of positively charged pions evaluated at mid-rapidity has a maximum at $\sqrt{s} \cong 4.5$ GeV and then decreases notably with the collision energy inside the region of $\sqrt{s} = 5 - 8$ GeV. This is shown in Fig. 1 *right*.



Figure 1: The mean transverse mass $\langle m_T \rangle - m_\pi$ of π^- (*left* panel) and π^+ (*right* panel) in $p+p$ reactions as a function of the center of mass energy $\sqrt{s}$. The rapidity window in the center of mass system is $0 < y < 0.2$. The solid lines show the results of UrQMD-3.4 calculations. The symbols with error bars in the *left* panel are the data from Ref. [16].

The nonmonotonous dependence of $\langle m_T \rangle$ on $\sqrt{s}$ seen in Fig. 1 *right* for π^+ accepted at mid-rapidity are absent for the UrQMD results taken at all rapidities. This is in an agreement with a compilation of the old $p+p$ data in Ref. [21].

A physical origin of the nonmonotonous dependence of $\langle m_T \rangle$ on $\sqrt{s}$ in the UrQMD model for π^+ in $p+p$ reactions is connected to a presence of two different sources of pions. Main inelastic reactions at small energies are the following [17, 18]:

$$p+p \to p+\Delta^+ \,, \quad p+p \to n+\Delta^{++} \,, \quad p+p \to \Delta^+ + \Delta^+ \,, \quad p+p \to \Delta^0 + \Delta^{++} \,,$$
$$p+p \to p+N^+ \,, \quad p+p \to N^+ + \Delta^+ \,, \quad p+p \to N^0 + \Delta^{++} \,. \tag{1}$$

In the UrQMD transport model, the states of N^* with $m = 1440, \ldots, 2250$ MeV and the states of Δ with $m = 1232, \ldots, 1950$ MeV are included. Specific charge states of N^* and Δ emerged in Eq. (1) explain a difference in a production of π^- and π^+: only Δ^0 and N^0 may produce π^- after their decays, whereas Δ^+ and N^+ may produce only π^+, and Δ^{++} has to produce π^+.

The reactions listed in (1) give the dominant contribution to the $p+p$ inelastic cross section at small collision energies. However, at $\sqrt{s} \geq 4$ GeV the excitations of baryonic strings,

$$p \ + \ p \ \rightarrow \ \text{String} \ + \ \text{String} \ , \tag{2}$$

open the new channels of hadron production. At collision energies $\sqrt{s} > 6$ GeV the string production dominates in the UrQMD description of inelastic $p+p$ cross section [17].

We can distinguish two sources contributing to hadron production in $p+p$ reactions: 1) the excitations of baryonic resonances listed in (1) and their decays to final hadrons; 2) the excitation of strings according to (2) and their decays to final hadrons. The mean transverse mass of a final hadron can be presented as

$$\langle m_T \rangle \ = \ f_B \, \langle m_T \rangle_B \ + \ f_S \, \langle m_T \rangle_S \ , \tag{3}$$

where $f_B(\sqrt{s})$ and $f_S(\sqrt{s})$ are the fractions of multiplicities for a given hadron species from excited baryons (1) and from strings (2), respectively. In Eq. (3), $\langle m_T \rangle_B$ and $\langle m_T \rangle_S$, also dependent on $\sqrt{s}$, are the values of the mean transverse mass for a hadron originating from these two sources. At SPS energies, the UrQMD gives $f_S > f_B$ for π^+. Besides, f_S increases and f_B decreases with $\sqrt{s}$. On the other hand, there is an opposite inequality, $\langle m_T \rangle_B > \langle m_T \rangle_S$, between the transverse masses of π^+ produced by baryons and by strings. This has a simple kinematic origin. To produce π^+ at mid-rapidity according to Eq. (1) baryonic resonances N^* or Δ should be themselves created in the mid-rapidity region. Thus, with increasing $\sqrt{s}$ transverse momentum and/or mass of the baryonic resonance has to increase. This leads to an increase of $\langle m_T \rangle_B$ of final π^+ with $\sqrt{s}$. An interplay of these two contributions to $\langle m_T \rangle$ in Eq. (3) leads to a nonmonotonous dependence on $\sqrt{s}$ of the resulting mean transverse mass as it seen in Fig. 1 *right* (see Ref. [15] for further details).

Note that K^- and $\overline{p}$ can not appear from decays of the baryonic resonances contributed to Eq. (1). In Fig. 2 *left* we present the UrQMD values of $\langle m_T \rangle - m$ at mid-rapidity for these hadrons as the functions of collision energy. No nonmonotonous dependence of $\langle m_T \rangle$ on $\sqrt{s}$ are seen.

Protons are obviously produced from decays of baryonic resonances excited according to Eq. (1). For heavy N^* the decay channel $N^* \rightarrow \Lambda + K^+$ is included in the UrQMD model.

Therefore, Eq. (3) can be also applied to a description of $\langle m_T \rangle$ for p, Λ, and K^+ in $p+p$ reactions. These UrQMD results are shown in Fig. 2 *right*. A nonmonotonous dependence of $\langle m_T \rangle - m$ on $\sqrt{s}$ at mid-rapidity ($0 < y < 0.2$) are observed for these hadron species. Like in a case of π^+, it disappears if final hadrons are detected in unrestricted rapidity window.



Figure 2: The results of the UrQMD-3.4 calculations for the mean transverse masses $\langle m_T \rangle - m$ in $p+p$ reactions as a function of collision energy. The rapidity window in the center of mass system is $0 < y < 0.2$. The error bars depict statistical uncertainties. *Left*: K^- (dashed line) and anti-protons (dotted line). *Right:* Protons (solid line), Λ (dashed line), and K^+ (dashed-dotted line).

A nonmonotonous energy dependence for the mid-rapidity values of $\langle m_T \rangle$ for π^+, p, K^+, and Λ in $p+p$ reactions observed in the UrQMD calculations is a consequence of an interplay of the contributions to the particle production from decays of the baryonic resonances and strings. Note that UrQMD includes all N^* and Δ states listed in the Particle Data Tables. A main difference between the baryonic resonances and strings consists of in rather different physical formulation of their decays to final hadrons. Note, however, that experimental information about the decay branching ratios of the high-mass resonance states is rather poor. In Figs. 3 *left* and *right* we present the results for π^+ and protons, respectively, obtained within two different versions of UrQMD, 3.3 and 3.4, as well as within the Hadron String Dynamics (HSD-2.5) [22–24]. The number of baryon species in the HSD model is smaller than that in UrQMD. In fact, not all baryonic resonances from the Particle Data Tables are included. Note that inelastic $p+p$ reactions above $\sqrt{s} \cong 2.6$ GeV are described in HSD-2.5 by the FRITIOF string model [25].



Figure 3: The mean transverse mass $\langle m_T \rangle - m$ of π^+ (*left* panel) and protons (*right* panel) in $p+p$ reactions as a function of collision energy. The rapidity window in the center of mass system is $0 < y < 0.2$. Solid lines correspond to the results of UrQMD-3.4, dashed ones to UrQMD-3.3, and dotted ones to HSD-2.5.

As seen from Fig. 3, both UrQMD-3.3 and UrQMD-3.4 predict a nonmonotonous dependence of $\langle m_T \rangle$ on $\sqrt{s}$ for π^+ and protons at mid-rapidity, whereas the HSD results give a monotonous increase of $\langle m_T \rangle$ with collision energy for both π^+ and protons.

In conclusion, within the UrQMD simulations we have observed a nonmonotonous dependence on the collision energy of the mean transverse mass $\langle m_T \rangle$ at mid-rapidity in inelastic $p+p$ collisions: a drop for π^+ at $5 \leq \sqrt{s} \leq 8$ GeV, a drop for K^+ at $4 \leq \sqrt{s} \leq 7$ GeV, a decrease for protons at $5 \leq \sqrt{s} \leq 17.3$ GeV and a decrease for Λ at $8 \leq \sqrt{s} \leq 17.3$ GeV. These results are the specific feature of UrQMD due to a presence of two mechanisms of the particle production: the decays of baryonic resonances, N^* and Δ, and strings. An interplay of these sources of particles at different collision energies results in nonmonotonous dependence of $\langle m_T \rangle$ on $\sqrt{s}$. Such a nonmonotonous dependence is not found in the HSD model, where the mass spectrum of N^* and Δ is underpopulated. Whether a nonmonotonous dependence of $\langle m_T \rangle$ on $\sqrt{s}$ at mid-rapidity for π^+, protons, K^+, and Λ is relevant for real $p+p$ interactions will be checked soon experimentally by the NA61/SHINE Collaboration. These experimental data can be used for evaluation of the unknown decay parameters of heavy resonances and inclusion of this information in the transport models.

Acknowledgments

We would like to thank Elena Bratkovskaya and Marek Gaździcki for fruitful discussions and comments. Publication is based on the research provided by the grant support of the State Fund for Fundamental Research (project No. F58/175-2014). The work was partially supported by the Helmholtz International Center for FAIR within the LOEWE program of the State of Hesse and by the Program of Fundamental Research of the Department of Physics and Astronomy of NAS of Ukraine (project No. 0112U000056).

---

[1] S.V. Afanasiev *et al.* [NA49 Collaboration], Phys. Rev. C 66 (2002) 054902.

[2] C. Alt *et al.* [NA49 Collaboration], Phys. Rev. C 73 (2006) 044910.

[3] C. Alt *et al.* [NA49 Collaboration], Phys. Rev. C 77 (2008) 024903.

[4] M. Gazdzicki and M. I. Gorenstein, Acta Phys. Polon. B 30 (1999) 2705.

[5] M. Gazdzicki, M. I. Gorenstein, and P. Seyboth, Acta Phys. Polon. B 42 (2011) 307.

[6] M. Gazdzicki, M. I. Gorenstein, and P. Seyboth, Int. Journ. Mod. Phys. E 23 (2014) 1430008.

[7] L. Van Hove, Phys. Lett. B 118 (1982) 138.

[8] C.M. Hung and E.V. Shuryak, Phys. Rev. Lett. 75 (1995) 4003.

[9] M. I. Gorenstein, M. Gazdzicki, and K. A. Bugaev, Phys. Lett. B 567 (2003) 175.

[10] M. Gaździcki [NA61/SHINE Collaboration], J. Phys. G 36 (2009) 064039.

[11] N. Abgrall *et al.* [NA61 Collaboration], JINST 9 (2014) P06005.

[12] G. S. F. Stephans, J. Phys. G 32 (2006) S447; G. Odyniec, PoS(CPOD 2013) 043.

[13] B. Friman, C. Hohne, J. Knoll, S. Leupold, J. Randrup, R. Rapp, and P. Senger, Lect. Notes Phys. 814 (2011) 1.

[14] P. Senger [CBM Collaboration], Cent. Eur. J. Phys., 10 (2012) 1289; see also http://www.fair-center.eu/for-users/experiments/cbm.html.

[15] V.Yu. Vovchenko, D.V. Anchishkin, and M.I. Gorenstein, Phys. Rev. C 90 (2014) 024916.

[16] N. Abgrall *et al.* [NA61 Collaboration], Eur. Phys. J. C 74 (2014) 2794.

[17] S.A. Bass *et al.*, Prog. Part. Nucl. Phys. 41 (1998) 255.

[18] M. Bleicher *et al.*, J. Phys. G 25 (1999) 1859.

[19] H. Petersen, M. Bleicher, S.A. Bass, and H. Stöcker, arXiv:0805.0567 [hep-ph].

[20] UrQMD-3.4 code is available at http://urqmd.org/

[21] A.M. Rossi *et al.*, Nucl. Phys. B 84 (1975) 269.

[22] W. Ehehalt and W. Cassing, Nucl. Phys. A 602 (1996) 449.

[23] J. Geiss, W. Cassing, and C. Greiner, Nucl. Phys. A 644 (1998) 107.

[24] W. Cassing and E.L. Bratkovskaya, Phys. Rept. 308 (1999) 65.

[25] B. Nilsson-Almqvist and E. Stenlund, Comput. Phys. Commun. 43 (1987) 387.